

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

WHAT IS CLAIMED IS:

- 5
1. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:
- (a) the nucleotide sequence as set forth in either SEQ ID NO: 1 or SEQ ID NO: 4;
- (b) the nucleotide sequence of the DNA insert in ATCC Deposit No. PTA-976;
- 10 (c) a nucleotide sequence encoding the polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;
- (d) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (c); and
- (e) a nucleotide sequence complementary to any of (a) - (c).
- 15 2. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:
- (a) a nucleotide sequence encoding a polypeptide which is at least about 70 percent identical to the polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, wherein the encoded polypeptide has an activity of the
- 20 polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;
- (b) a nucleotide sequence encoding an allelic variant or splice variant of the nucleotide sequence as set forth in either SEQ ID NO: 1 or SEQ ID NO: 4, the nucleotide sequence of the DNA insert in ATCC Deposit No. PTA-976, or (a);
- (c) a region of the nucleotide sequence of either SEQ ID NO: 1 or
- 25 SEQ ID NO: 4, the DNA insert in ATCC Deposit No. PTA-976, (a), or (b) encoding a polypeptide fragment of at least about 25 amino acid residues, wherein the polypeptide fragment has an activity of the encoded polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, or is antigenic;
- (d) a region of the nucleotide sequence of either SEQ ID NO: 1 or
- 30 SEQ ID NO: 4, the DNA insert in ATCC Deposit No. PTA-976, or any of (a) - (c) comprising a fragment of at least about 16 nucleotides;

(e) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (d); and

(f) a nucleotide sequence complementary to any of (a) - (d).

5 3. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one conservative amino acid substitution, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

(b) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid insertion, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

15 (c) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid deletion, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

(d) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 which has a C- and/or N- terminal truncation, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

20 (e) a nucleotide sequence encoding a polypeptide as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the encoded polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

25 (f) a nucleotide sequence of any of (a) - (e) comprising a fragment of at least about 16 nucleotides;

30

- Sul*
A
- (g) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (f); and
(h) a nucleotide sequence complementary to any of (a) - (e).

5 4. A vector comprising the nucleic acid molecule of any of Claims 1, 2, or 3.

5. A host cell comprising the vector of Claim 4.

Sul
C 10 6. The host cell of Claim 5 that is a eukaryotic cell.

7. The host cell of Claim 5 that is a prokaryotic cell.

15 8. A process of producing an IFN-L polypeptide comprising culturing the host cell of Claim 5 under suitable conditions to express the polypeptide, and optionally isolating the polypeptide from the culture.

9. A polypeptide produced by the process of Claim 8.

20 10. The process of Claim 8, wherein the nucleic acid molecule comprises promoter DNA other than the promoter DNA for the native IFN-L polypeptide operatively linked to the DNA encoding the IFN-L polypeptide.

25 11. The isolated nucleic acid molecule according to Claim 2, wherein the percent identity is determined using a computer program selected from the group consisting of GAP, BLASTN, FASTA, BLASTA, BLASTX, BestFit, and the Smith-Waterman algorithm.

30 12. A process for determining whether a compound inhibits IFN-L polypeptide activity or IFN-L polypeptide production comprising exposing a cell

according to any of Claims 5, 6, or 7 to the compound and measuring IFN-L polypeptide activity or IFN-L polypeptide production in said cell.

5 13. An isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5; and

(b) the amino acid sequence encoded by the DNA insert in ATCC Deposit No. PTA-976.

10

14. An isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in either SEQ ID NO: 3 or SEQ ID NO: 6, optionally further comprising an amino-terminal methionine;

15 (b) an amino acid sequence for an ortholog of either SEQ ID NO: 2 or SEQ ID NO: 5;

(c) an amino acid sequence which is at least about 70 percent identical to the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

20

(d) a fragment of the amino acid sequence set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 comprising at least about 25 amino acid residues, wherein the fragment has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, or is antigenic; and

25 (e) an amino acid sequence for an allelic variant or splice variant of the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, the amino acid sequence encoded by the DNA insert in ATCC Deposit No. PTA-976, or any of (a) - (c).

30

15. An isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one conservative amino acid substitution, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

5 (b) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid insertion, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

(c) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one amino acid deletion, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5;

(d) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 which has a C- and/or N- terminal truncation, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5; and

15 (e) the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5.

20

16. An isolated polypeptide encoded by the nucleic acid molecule of any of Claims 1, 2, or 3, wherein the polypeptide has an activity of the polypeptide set forth in either SEQ ID NO: 2 or SEQ ID NO: 5.

25

17. The isolated polypeptide according to Claim 14, wherein the percent identity is determined using a computer program selected from the group consisting of GAP, BLASTP, FASTA, BLASTA, BLASTX, BestFit, and the Smith-Waterman algorithm.

30

18. A selective binding agent or fragment thereof which specifically binds the polypeptide of any of Claims 13, 14, or 15.

19. The selective binding agent or fragment thereof of Claim 18 that specifically binds the polypeptide comprising the amino acid sequence as set forth in either SEQ ID NO: 2 or SEQ ID NO: 5, or a fragment thereof.

5

20. The selective binding agent of Claim 18 that is an antibody or fragment thereof.

21. The selective binding agent of Claim 18 that is a humanized antibody.

10

22. The selective binding agent of Claim 18 that is a human antibody or fragment thereof.

15

23. The selective binding agent of Claim 18 that is a polyclonal antibody or fragment thereof.

20

24. The selective binding agent Claim 18 that is a monoclonal antibody or fragment thereof.

25. The selective binding agent of Claim 18 that is a chimeric antibody or fragment thereof.

25

26. The selective binding agent of Claim 18 that is a CDR-grafted antibody or fragment thereof.

27. The selective binding agent of Claim 18 that is an antiidiotypic antibody or fragment thereof.

30

28. The selective binding agent of Claim 18 that is a variable region fragment.

29. The variable region fragment of Claim 28 that is a Fab or a Fab' fragment.

5 30. A selective binding agent or fragment thereof comprising at least one complementarity determining region with specificity for a polypeptide having the amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5.

31. The selective binding agent of Claim 18 that is bound to a
10 detectable label.

32. The selective binding agent of Claim 18 that antagonizes IFN-L polypeptide biological activity.

15 33. A method for treating, preventing, or ameliorating an IFN-L polypeptide-related disease, condition, or disorder comprising administering to a patient an effective amount of a selective binding agent according to Claim 18.

34. A selective binding agent produced by immunizing an animal with
20 a polypeptide comprising an amino acid sequence of either SEQ ID NO: 2 or SEQ ID NO: 5.

35. A hybridoma which produces a selective binding agent which is
25 capable of binding a polypeptide according to any of Claims 1, 2, or 3.

36. A method of detecting or quantitating the amount of IFN-L polypeptide using the anti-IFN-L antibody or fragment of Claim 18.

37. A composition comprising the polypeptide of any of Claims 13,
30 14, or 15, and a pharmaceutically acceptable formulation agent.

39. The composition of Claim 37, wherein the polypeptide comprises the amino acid sequence as set forth in either SEQ ID NO: 3 or SEQ ID NO: 6.

10

20

25

30

46. A fusion polypeptide comprising the polypeptide of any of Claims 13, 14, or 15 fused to a heterologous amino acid sequence.

Jul
C1

10

15

20

25

51. A device, comprising:

(a) a membrane suitable for implantation; and

(b) cells encapsulated within said membrane, wherein said cells secrete a protein of any of Claims 13, 14, or 15; and

said membrane is permeable to said protein and impermeable to materials detrimental to said cells.

52. A method of identifying a compound which binds to an IFN-L
30 polypeptide comprising:

(a) contacting the polypeptide of any of Claims 13, 14, or 15 with a compound; and

(b) determining the extent of binding of the IFN-L polypeptide to the compound.

5

53. The method of Claim 52, further comprising determining the activity of the polypeptide when bound to the compound.

54. A method of modulating levels of a polypeptide in an animal comprising administering to the animal the nucleic acid molecule of any of Claims 1, 2, or 3.

55. A transgenic non-human mammal comprising the nucleic acid molecule of any of Claims 1, 2, or 3.

15

56. A process for determining whether a compound inhibits IFN-L polypeptide activity or IFN-L polypeptide production comprising exposing a transgenic mammal according to Claim 54 to the compound, and measuring IFN-L polypeptide activity or IFN-L polypeptide production in said mammal.